

## WOUND HEALING ACTIVITY OF ALCOHOLIC EXTRACT OF *KAEMPFERIA GALANGA* IN WISTAR RATS

SHANBHAG TARA V., SHARMA CHANDRAKALA,  
ADIGA SACHIDANANDA, BAIRY LAXMINARAYANA KURADY\*,  
SHENOY SMITA AND SHENOY GANESH

*Department of Pharmacology,  
Kasturba Medical College,  
Manipal – 576 104*

( Received on September 6, 2005 )

**Abstract :** The wound healing effect of alcoholic extract of *Kaempferia galanga* (*K. galanga*) and its effect in dexamethasone suppressed wound healing was studied in Wistar rats. Three wound models viz. incision, excision and dead space wounds were used in this study. The parameters studied were breaking strength in case of incision wounds, epithelialization and wound contraction in case of excision wound and granulation tissue dry weight, breaking strength and hydroxyproline content in case of dead space wound. The dexamethasone treated group showed a significant ( $P < 0.001$ ) reduction in the wound breaking strength when compared to control group in incision type of wound model. Co-administration of *K. galanga* with dexamethasone had significantly ( $P < 0.001$ ) increased the breaking strength of dexamethasone treated group. In excision wound model, the percentage of the wound contraction was significantly ( $P < 0.05$ ) increased by *K. galanga* only on 16th day and also it reversed the dexamethasone suppressed wound contraction on the 16 day. *K. galanga* significantly ( $P < 0.001$ ) reduced the time required for epithelialization and reversed the epithelialization delaying effect of dexamethasone significantly ( $P < 0.001$ ).

**Key words :** *Kaempferia galanga* dexamethasone  
wound contraction wound breaking strength  
period of epithelialization

### INTRODUCTION

Wound is a breach in the normal tissue continuum, resulting in a variety of cellular and molecular sequelae. The basic principles of optimal wound healing which include minimizing tissue damage, debriding non-viable tissue, maximizing tissue perfusion

and oxygenation, proper nutrition and moist wound healing environment have been recognized for many years (1). A number of drugs ranging from simple non-expensive analgesics to complex and expensive chemotherapeutic agents administered in the management of wound affect healing either positively or negatively (2). Aspirin,

---

\*Corresponding Author : E-mail : klbairy@yahoo.com

indomethacin, cytotoxic agents and immunosuppressant have been proved experimentally to affect healing negatively (3, 4, 5, 6).

Medicinal herbs are an indispensable part of traditional medicine. The rhizome of *K. galanga* finds an important place in indigenous medicine as an expectorant, diuretic and carminative (7). It is also found to have anticancer (8), antihypertensive (9) and larvicidal activity (10). It is used for the treatment of various skin disorders, rheumatism and diabetes mellitus (11, 12). However to the best of our knowledge a systematic study on wound healing activity of *K. galanga* has not been undertaken. Hence, the present study was undertaken to evaluate the wound healing property of alcoholic extract of *K. galanga* rhizome and to study its influence on dexamethasone suppressed wound healing on various animal wound models in Wistar rats.

## MATERIALS AND METHODS

### Collection and preparation of alcoholic extract of *Kaempferia galanga*:

*K. galanga* plants were procured locally in the month of December and authenticated by Professor of Botany, Mahatma Gandhi Memorial College, Udupi. The shade dried rhizomes were crushed into small pieces and powdered. The powder was loaded into soxhlet extractor in 8 batches of 250 g each and was subjected to extraction for about 30–40 h with ethanol 95%. After extraction the solvent was distilled off and the extract was concentrated under reduced pressure on a water bath at a temperature below 50°C to a syrupy consistency. Then it was

dried in the dessicator. The yield was about 3%.

### Animal care and Handling:

This was done as per the guidelines set by the Indian National Science Academy New Delhi, India. Twelve-week-old healthy Wistar rats (150–200 g) of either sex bred locally in the animal house of Kasturba Medical College, Manipal were selected for the study. They were housed under controlled conditions of temperature of  $23 \pm 2^\circ\text{C}$ , humidity of  $50 \pm 5\%$  and 10–14 h of light and dark cycles respectively. The animals were housed individually in polypropylene cages containing sterile paddy husk (procured locally) as bedding throughout the experiment and had free access to sterile food (animal chow) (M/s Hindustan Lever Ltd.) and water *ad libitum*. The study was undertaken after obtaining the approval of Institutional Animal Ethical Committee (IEAC approval letter No. IEAC/KMC/04/2002–2003 Dated Dec. 30th, 2003).

### Study Design:

The animal were randomly allocated into four groups of eight animals each for the three experimental animal wound models.

Group I received 2 ml of gum acacia 2% (E. Merck India Ltd.) po through intragastric tube.

Group II received *K. galanga*, 300 mg/kg po. The dose selection was based on the toxicity studies.

Group III received dexamethasone, 0.17 mg/kg (13) (Cadila Healthcare, Mumbai) im.

Group IV received dexamethasone (0.17 mg/kg im) & *K. galanga* (300 mg/kg) po.

The suspension of the alcoholic extract of *K. galanga* was made in 2% gum acacia. The dose selection was based on the toxicity studies. In group IV, extract of *K. galanga* was administered immediately after intramuscular injection of dexamethasone.

**Acute Toxicity Studies:** Healthy Wistar rats of either sex were chosen and were divided into four groups (n=6). They were administered single dose of alcoholic extract of *K. galanga* orally with increasing doses of 100, 300, 1000, 3000 mg/kg body weight respectively. The doses upto 3 g/kg was well tolerated without producing any signs of toxicity and mortality. 10% of the maximum tolerated dose i.e. 300 mg/kg was selected for the study.

**Dosing Schedule:** *K. galanga* extract and dexamethasone were administered orally and intramuscularly respectively once daily from day 0 to day 9 in the incision and dead space wound models and from day 0 to the day of complete healing or the 21st postoperative day, whichever occurred earlier in the excision wound model. In group IV *K. galanga* extract was given after the injection of dexamethasone.

#### Wound models

All wounding procedures were carried out under pentobarbitone (Rhone-poulenc B.P., France) (3 mg/100 g) anesthesia. In the present study no animal showed visible signs of infection.

**1. Incision wound:** On the depilated backs of the animals, two paravertebral incisions of 6 cm length were made cutting through the full thickness of the skin. Interrupted sutures, 1 cm apart, were placed to approximate the cut edges of the skin (14). The sutures were removed on the 7th post wound day and skin breaking strength was measured on the 10th day by continuous water flow technique of Lee (3).

**2. Dead space wound:** Dead space wounds were created through a small transverse incision made in the lumbar region (15). A polypropylene tube (2.5 × 0.5 cm) was implanted subcutaneously beneath the dorsal paravertebral lumbar skin. The day of the wound creation was considered as day zero. Granulation tissue formed on the polypropylene tube was harvested by careful dissection on day 10 and the breaking strength of the granulation tissue was measured. The granulation tissue was dried in an oven at 60°C overnight and the dry weight was noted. Acid hydrosylate of the dry tissue was used for the determination of the hydroxyproline content (16).

**2. Excision wound:** An excision wound was inflicted by cutting away 500 mm<sup>2</sup> full thickness of a pre-determined area on the depilated back of the rat. Epithelialization period was noted as the number of days after wounding required for the eschar to fall off leaving no raw wound behind. Wound contraction rate was monitored by planimetric measurement of the wound area on alternate days. This was achieved by tracing the wound on a graph paper. Reduction in the wound area was expressed as percentage of the original wound size (17).

**Statistical analysis**

Results were analysed by One way analysis of variance (ANOVA) followed by Scheffe's test using SPSS computer package version-11.

**RESULTS****Incision wound model**

The mean breaking strength in the control group was  $348.27 \pm 7.8$  g. The alcoholic extract of *K. galanga* did not alter the breaking strength when compared to control. In the dexamethasone treated group the mean breaking strength was  $166.03 \pm 7.45$  g which was significantly ( $P < 0.001$ ) less compared to control group. Coadministration of *K. galanga* with dexamethasone has significantly ( $P < 0.001$ ) increased the breaking strength to  $292.6 \pm 11.72$  g (Table I).

**Dead space wound model**

The mean breaking strength of granulation tissue in the control group was  $263.75 \pm 28.59$  g. Even though there was no significant alteration in the breaking strength of the granulation tissue, a marked increase in breaking strength was observed ( $312.5 \pm 37.4$  g) in *K. galanga* treated group when compared to the control group. The breaking strength in dexamethasone treated group was  $273.75 \pm 12.09$  g. The increase in the breaking strength compared to the control group can not be explained. (Table I). The mean dry weight of granulation tissue in control group was  $42.12 \pm 5.47$  mg which was significantly ( $P < 0.05$ ) increased to  $49.75 \pm 5.56$ ,  $64.00 \pm 6.81$ ,  $61.87 \pm 6.15$  mg in groups treated with *K. galanga*, dexamethasone, dexamethasone + *K. galanga* respectively (Table I). The increase in dry weight in dexamethasone group could be due to fibroblasts and other inflammatory cells.

TABLE I: Effect of *K. galanga* on dead space and incision wound parameters.

Drugs (n=8)	Dose/route	Dead space wound			Incision wound
		Breaking strength of granulation tissue (g) Mean $\pm$ S.E.	Dry weight of granulation tissue – mg Mean $\pm$ S.E.	Hydroxyproline content – mg/g of the tissue Mean $\pm$ S.E.	Breaking strength (g) Mean $\pm$ S.E.
Gum acacia	2 ml oral	263.75 $\pm$ 28.59	42.12 $\pm$ 5.47	21.09 $\pm$ 4.41	348.27 $\pm$ 7.8
<i>K. galanga</i>	300 mg/kg oral	312.5 $\pm$ 37.4	49.75 $\pm$ 5.56 <sup>b</sup>	29.61 $\pm$ 4.88	349.78 $\pm$ 9.13
Dexa	0.17 mg/kg im	273.75 $\pm$ 12.09	64 $\pm$ 6.81 <sup>b</sup>	19.85 $\pm$ 2.69	166.03 $\pm$ 7.45 <sup>c</sup>
Dexa+ <i>K. galanga</i>	0.17 mg/kg im + 300 mg/kg oral	386.25 $\pm$ 10.34 <sup>a</sup>	61.87 $\pm$ 6.15 <sup>b</sup>	23.14 $\pm$ 2.50	292.6 $\pm$ 11.72 <sup>d</sup>

Dexa=Dexamethasone.

<sup>a</sup>P<0.05 Vs Dexamethasone, Oneway ANOVA, F=5.004, df=3, 28.

<sup>b</sup>P<0.05 Vs Control, Oneway ANOVA, F=2.939, df=3, 28.

<sup>c</sup>P<0.001 Vs Control, Oneway ANOVA, F=88.249, df=3, 28.

<sup>d</sup>P<0.001 Vs Dexamethasone, Oneway ANOVA, F=88.249, df=3, 28.

TABLE II: Effect of *K. galanga* on excision wound parameters.

Drugs	Dose/route	Percentage of wound contraction (Mean±S.E.)				Period of epithelialization (days) Mean±S.E.
		4th day	8th day	12th day	16th day	
G. acacia	2 ml oral	27.75±4.38	47.15±5.25	59.45±2.77	68.67±1.28	16.75±0.75
<i>K. galanga</i>	300 mg/kg oral	21.2±3.21	48.25±4.46	67.55±3.48	82.1±2.22 <sup>a</sup>	11.12±0.47 <sup>c</sup>
Dexa	0.17 mg/kg im	23.4±3.32	39.57±3.58	55.85±2.39	67.02±2.12	17.25±0.75
Dexa+ <i>K. galanga</i>	0.17 mg/kg im + 300 mg/kg oral	26.25±3.32	37.5±2.64	65.77±0.93	76.22±1.03 <sup>b</sup>	12.75±0.77 <sup>d</sup>

Dexa=Dexamethasone.

<sup>a</sup>P<0.001 Vs Control, Oneway ANOVA, F=5.004, df=3, 28.

<sup>b</sup>P<0.001 Vs Dexamethasone, Oneway ANOVA, F=2.939, df=3, 28.

<sup>c</sup>P<0.001 Vs Control, Oneway ANOVA, F=18.483, df=3, 28.

<sup>d</sup>P<0.001 Vs Dexamethasone, Oneway ANOVA, F=18.483, df=3, 28.

The mean hydroxyproline content of granulation tissue in control group was  $21.09 \pm 4.41$  mg/g of the tissue. It was not significantly altered in any of the groups (Table I).

#### Excision wound

The percentage of wound contraction was  $27.75 \pm 4.38$ ,  $47.15 \pm 5.25$ ,  $59.45 \pm 2.77$  and  $68.67 \pm 1.28$  as measured on the 4th, 8th, 12th and 16th day respectively in the control group. The wound contraction rate was not altered significantly in any of the test groups on 4th, 8th and 12th day as compared to control group at same time. Apart from this, we have also noted a positive trend in wound contraction rate in *K. galanga* treated group and negative trend in wound contraction rate in dexamethasone treated group even though they were not statistically significant on 8th and 12th day. However wound contraction-rate was significantly increased in *K. galanga* treated group compared to the control group on 16th day (P<.001) ( $82.1 \pm 2.22$ ). Similar observation was also made in the

dexamethasone & *K. galanga* treated group when compared to the dexamethasone treated group where it increased from  $67.02 \pm 2.12$  to  $76.22 \pm 1.03$  on 16th day (P<0.001) (Table II). The mean period of epithelialization in the control group was  $16.75 \pm 0.75$  days. It was significantly (P<.001) reduced to  $11.12 \pm 0.47$  days in *K. galanga* treated group. The mean period of epithelialization in dexamethasone treated group was  $17.25 \pm 0.75$  days which was significantly (P<0.001) reduced to  $12.75 \pm 0.77$  days in the group treated with both dexamethasone and *K. galanga* (Table II).

#### DISCUSSION

Granulation, collagen maturation and scar formation are some of the many phases of wound healing which run concurrently, but independent of each other. The use of single model is inadequate and no reference standard exists that can collectively represent the various phases of wound healing. Hence three different models have

been chosen in our study to assess the effect of *K. galanga* on wound healing. The wound breaking strength is determined by the rate of collagen synthesis and more so by the maturation process where there is covalent binding of collagen fibrils through inter and intra molecular cross linking. In our study dead space wound model showed no significant increase in breaking strength and hydroxyproline concentration, but the dry weight of the granulation tissue was significantly increased in *K. galanga* treated group. By this we can assume that the *K. galanga* might not have increased the collagen content but probably have altered the maturation process, by affecting the cross linking of collagen or improving the quality of collagen fibrils. The increase in weight in dexamethasone treated group could be due to high protein concentration and collagen bundle formation (18). It is difficult to explain the effect of *K. galanga* along with the dexamethasone as there was a slight increase in breaking strength and dry weight of granulation tissue in the dexamethasone alone treated group compared to control group.

Wound contraction is the process of mobilizing healthy skin surrounding the wound to cover the denuded area. This centripetal movement of wound margin is believed to be due to the activity of myofibroblast (19). Since *K. galanga* enhanced wound contraction, it would have either enhanced contractile property of myofibroblasts or increased the number of myofibroblasts recruited into the wound area. In excision wound model *K. galanga* hastened the period of epithelialization

significantly and the co-administration of *K. galanga* with dexamethasone hastened the epithelialization in dexamethasone group. Even though only during later part, *K. galanga* showed significant increase in wound contraction we have observed the positive trend in the initial stages. Concomitant administration of *K. galanga* along with dexamethasone had also significantly increased the wound contraction on 16th day. Hence it appears that *K. galanga* has prohealing effect as evidenced by the above findings. It also appears that *K. galanga* was able to promote epithelialization either by facilitating the proliferation of epithelial cells or by increasing the viability of epithelial cells. It is difficult to draw any conclusion from the study regarding the dexamethasone & *K. galanga* effect in dexamethasone suppressed wound model.

In recent years oxidative stress has been implicated in a variety of degenerative process and diseases. These include acute and chronic inflammatory condition such as wound healing (20). *K. galanga* has shown to possess anti-oxidant property (21). The flavonoids which are responsible for the free radical scavenging activity were believed to be one of the important components in wound healing (22). Phytochemical screening revealed the presence of flavonoids in *K. galanga* (23). This could be the reason for prohealing activity of *K. galanga*. This enhanced wound contraction effect of *K. galanga* and epithelialization could possibly be made use of clinically in healing of open wounds. However confirmation of this suggestion will need well designed clinical evaluation.

## REFERENCES

1. Pierce GF, Mustoe TA. Pharmacologic enhancement of wound healing. *Annu Rev Med* 1995; 46: 467–481.
2. Prasad D, Rao CM. Wound healing profile of ketorolac, metronidazole and tinidazole administered postsurgically. *Ind J Exp Biol* 1995; 33: 845–847.
3. Lee KH. Studies on the mechanism of action of salicylates III. Effect of vitamin A on the wound healing retardation action of aspirin. *J Pharma Sci* 1968 July; 57(7): 1238–1240.
4. Rao CM, Ramesh KV, Bairy KL, Kulkarni DR. A simple method to quantify maturation of wound collagen. *Ind J Exp Biol* 1991; 29: 156–158.
5. Raju S, Kulkarni DR. Vitamin A reverses the wound healing suppressant effect of cyclophosphamide. *Ind J Pharmacol* 1986; 18: 154–157.
6. Holla RK, Sequeira RP, Kulkarni DR. Cyclosporin and wound healing. *Ind J Exp Biol* 1988; 26: 869–873.
7. Achuthan CR, Jose Padikkala. Hypolipidemic effect of *Alpinia galanga* (Rasna) and *Kaempferia galanga* (Kachoori). *Ind J Clin Biochem* 1997; 12(1): 55–58.
8. Xue Y, Chen H. Study on the anti-carcinogenic effect of three compounds of *Kaempferia galanga* L. *Xei Shen Yan Jiu* 2002; 31(4): 247–248, 251.
9. Othman R, Ibrahim H, Mohd MA, Awang K, Gilani AV, Musthafa MR. Vaso relaxant effect of *Kaempferia galanga* on smooth muscles of rat aorta. *Planta Med* 2002; Jul 68(7): 655–657.
10. Choochote W, Kanjanapothi D, Panthanga A, Taesotikul T, Jitpakdi A, Chaithong U, Pitasawat B. Larvicidal and repellent effects of *Kaempferia galanga*. *Southeast Asian J Trop Med Public Health* 1999; Sep 30(3): 470–476.
11. Mangaly JK, Sabu M. Ethanobotany of zingiberaceae. Zingiberaceae workshop. Prince of Songkla University, Hat Yai, Thailand. 1991; 15–18 Oct; p. 24.
12. Vagbhata. *Astanga Hridayam Chikitsa Stanum* (Saranga Sundara Commentary of Arun Datta and Hemadri) 3rd Chapter. Chaukhambha Orientalia, Varanasi, Delhi 1971; 167–681.
13. Shanbhag T, Shenoy S, Rao MC. Wound healing profile of *Tinospora cordifolia*. *Indian Drugs* 2005; 42(4): 217–221.
14. Lee KH, Tong TG. Mechanism of action of retinyl compounds on wound healing II. Effect of active retinyl derivative on granuloma formation. *J Pharma Sci* 1970; 59: 1195–1197.
15. Nayak S, Rao SG, Murthy KD, Somayaji SN, Bairy KL. Pyramid environment reduces the wound healing suppressant properties of dexamethasone in albino rat. *Indian J Expt Biol* 2003; Jun 41(6): 645–648.
16. Keuman RE, Logan MA. The determination of collagen and elastin in tissues. *J Biochem* 1972; 186: 549–556.
17. Bairy KL, Rao CM. Wound healing profile of *Ginkgo biloba*. *J Natural Remedies* 2001; 1: 25–27.
18. Annie Shirweiker, Radhika Shenoy, Udupa AL, Udupa SL, Somashekhar Shetty. Wound healing property of leaves of *Hyptis suaveolens* with supportive role of antioxidant enzymes. *Indian J Exp Biol* 2003; 41: 238–241.
19. Gabbaiani G, Harschel BJ, Ryan GB. Granulation tissue as a contractile organ. *J Exp Med* 1976; 135: 719.
20. Maiere CM, Chan ph. Role of superoxide dismutase in oxidative damage and neurodegenerative disorders. *Neuroscientist* 2002; 8: 323–324.
21. Jitoe A, Masuda T, Tengah IGP, Suprapta DN, Gara IW, Nakatani N. Antioxidant activity of tropical ginger extract and analysis of the contained curcuminoids. *J Agric Fd Che* 1992; 40(8): 1337–1340.
22. Devipriya S, Shyamaladevi CS. Protective effect of quercetin induced cell injury in the kidney. *Indian J Pharmacol* 1999; 422–424.
23. Truntiwachwuttikul P. The chemistry of *Kaempferia*. *Zingiberaceae workshop*. Prince of Sonkla University, Hat Yai, Thailand 1991, 15–18 Oct; p. 10.