https://ijpp.com





Short Communication

Indian Journal of Physiology and Pharmacology



Article in Press

Handgrip strength cut off value estimation in Indian older adults using LASI-1 dataset

Sambaran Mondal¹, Sudipta Banerjee¹, Shankarashis Mukherjee¹

¹Department of Food and Nutrition, West Bengal State University, West Bengal, Kolkata, India.

*Corresponding author:

Shankarashis Mukherjee, PhD, Professor and Head, Department of Food and Nutrition, West Bengal State University, Kolkata, West Bengal, India.

msasish@yahoo.co.in

Received: 17 October 2024 Accepted: 02 April 2025 EPub Ahead of Print: 09 June 2025 Published:

DOI 10.25259/IJPP_585_2024

Quick Response Code:



ABSTRACT

Objectives: Handgrip strength (HGS) is a well established indicator of sarcopenia. The study tried to report the distribution of HGS among the Indian older adults with a large dataset of LASI-1 and attempted to generate age and sex specified cut-off values following the procedure recommended by Asian Working Group for Sarcopenia.

Materials and Methods: After considering inclusion and exclusion criteria, 40351 healthy individuals aged 45-89 years of both sexes were selected. Arithmetic mean and standard deviation for HGS were found out. HGS cut-off values were generated using the lowest 20th percentile from the study population.

Results: Average HGS was found to be higher in females $(21.74 \pm 7.447 \text{ kg})$ than males $(21.67 \pm 7.361 \text{ kg})$. HGS cut-off value of 15.25 kg was generated for Indian older adults of both sexes.

Conclusion: Generated HGS cut-off value of 15.25 kg may be found to effective in early screening of sarcopenia among the Indian older adults.

Keywords: Handgrip strength, HGS cut-off, Indian older adults, sarcopenia

INTRODUCTION

Sarcopenia can be defined as a continuous and generalised degeneration of muscle strength and skeletal muscle mass, which can be identified through the measurement of handgrip strength (HGS).^[1] Sarcopenia can be categorised as primary sarcopenia, which is associated with ageing and secondary sarcopenia, which results from chronic conditions and diseases such as cancer, arthritis and so on.^[2]

Grip strength is an easy to administer and reliable measurement of maximum voluntary muscle contraction and also serves as an indicator of the overall muscular strength of the human body.^[3] It not only helps in predicting the muscle mass in the body but also provides an indication about the state of nutrition, the occurrence of chronic diseases, quality of life and mortality.^[4-6] Nutritional status plays a crucial role in maintaining optimum HGS by regulating the skeletal muscle mass in the body and any fluctuation in it, in terms of malnutritional state can seriously hamper the HGS and leads to early development of sarcopenia and frailty among the older adults.^[7,8]

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, transform, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms. ©2025 Published by Scientific Scholar on behalf of Indian Journal of Physiology and Pharmacology

In recent times, HGS has been widely used as a clinical indicator of deprived mobility and low muscle mass.^[9] HGS indicates the presence and function of muscle fibres in the body, which is affected by the distribution of muscle mass, and also helps to identify the early signs of sarcopenia in terms of deprived HGS, especially among older adults and the elderly.^[10]

The European Working Group on Sarcopenia in Older People (EWGSOP2) revised the older definitions (EWGSOP) and terms and set new cutoff values of HGS for indication of lower strength as <27 kg for men and <16 kg for women.^[11]As there is a disparity between European and Asian populations, in terms of ethnicity, body structure, climate and other related factors, the Asian Working Group for Sarcopenia (AWGS) proposed a HGS cutoff of <28 kg for men and <18 kg for women, which indicates lower strength.^[12] However, there is no standardised and specified cutoff values for the Indians.

In Indian context, the Longitudinal Aging Study in India (LASI), Wave 1 depicts a larger health scenario of the older adults and the elderly prevailing in the country in its report; but still there is a dearth of knowledge regarding the sarcopenia and its allied issues.^[13] Few attempts were made to specify cutoff values of HGS for Indian older adults, but most of them lack large sample sizes from the population.^[14,15] In this backdrop, the purpose of the study is to report the distribution of HGS among the Indian older adults and the elderly with age and sex specification and also generate HGS cutoff values for identification of lower muscle strength among them.

MATERIALS AND METHODS

Data

The present study used the dataset of LASI-1, which was conducted nationwide including the Union territories, during 2017-2018 with a multistage, stratified, cluster and area probabilistic random sampling method. The study consisted of 72250 individuals aged 45 years and above of both sexes. Data on several health parameters including the anthropometric assessments along with psychological and social health related parameters were collected using standard procedures and validated questionnaires. Grip strength was measured by LASI, using a handheld dynamometer (Smedley's Hand Dynamometer) of both hands (considering the dominant and non-dominant hands).^[13] The subjects were asked to sit in a chair with armrests to eliminate the influence of gravitational force, and instructed to press the dynamometer as hard as they could for a couple of seconds and then let go, 3 times in both hands and the maximum values were chosen as HGS; the trained health investigators collected two reading of HGS (nearest to 0.5 kg) for both dominant and non-dominant hands performed with a rest of 30 s between the two measurements.^[16] The present study used the calculated average HGS values for analysis.^[17]

Inclusion criteria

From the 72250 individual data, the present study included the data of 40,351 healthy individuals aged between 45 and 89 years of both sexes the other criterion being availability of data on grip strength of both hands.

Exclusion criteria

Individuals aged <45 years and \geq 90 years were excluded and excluded were those whose data for grip strength of both hands were not available. Individuals suffering from chronic diseases such as Alzheimer's, bone, or joint diseases such as arthritis, osteoporosis and rheumatism, diabetes, lung diseases, heart diseases, stroke and neurological or psychiatric problems were also excluded from the study.

HGS cutoff values for determination of sarcopenia

The EWGSOP recommended for defining the cutoff of HGS, a study should be conducted using the normative population, that is healthy young adults rather than other provided predictive reference populations and the cutoff points will be below 2 standard deviations (SDs) from the mean reference value.^[18] Another approach by AWGS stated that, when normative data is unavailable, cutoff values can be generated using the lowest quartiles or quintiles with sex-wise specification.^[19]

In the Indian context, in the absence of normative data, the study selected a healthy not so aged population and estimate normative data as well as used age and sex specific lowest 20th percentile to generate cutoff values of HGS for older Indian population.

Statistical analysis

All statistical analyses were performed using IBM SPSS Gradepack 29. Average strength (kg) of both hands was calculated. The individuals were categorised sex-wise as men and women and also specified with age groups as 45–49 years, 50–54 years, 55–59 years, 60–64 years, 65–69 years, 70–74 years, 75–79 years, 80–84 years and 85–89 years. Descriptive statistics such as mean and SDs were calculated for all variables. Percentiles of HGS were calculated using HAVERAGE method. The HGS cutoff values for older Indian population were generated using the age- and sex-adjusted lowest 20% quintiles in the study population.

RESULTS

Among the 40351 healthy older Indian individuals, 18833 (46.67%) were males and 21518 (53.33%) were females. The mean age of the males was 59.8 \pm 10.36 years (presented in arithmetic mean \pm standard deviation form) and 59.0 \pm 10.15 years for females. The anthropometric

Table 1: Anthropometric profile of the healthy individuals included in the study.									
Age groups (years)	Sex	n	Anthropometric parameters						
			BH (cm)	BW (cm)	BMI	WC (cm)			
45-49	Male	3787	155.1±8.73	54.6±12.75	22.60 ± 4.561	84.2±12.49			
	Female	4568	155.4±8.65	54.2±12.30	22.39 ± 4.438	83.7±12.34			
50-54	Male	3215	155.5±8.77	54.3±12.53	22.35 ± 4.450	83.8±12.49			
	Female	3672	155.2±8.76	54.1±12.26	22.43 ± 4.928	83.7±12.34			
55–59	Male	2765	154.9±8.53	53.8±12.36	22.32±4.483	83.6±12.24			
	Female	3362	155.1±8.83	53.9±12.15	22.36±4.707	83.6±12.10			
60–64	Male	2929	155.3±8.65	$54.0{\pm}12.48$	22.31±4.452	83.5±12.48			
	Female	3456	155.4±8.61	54.5±12.77	22.46±4.616	84.0±12.58			
65–69	Male	2658	155.1±8.87	54.5±12.65	22.57±4.636	84.0±12.21			
	Female	2765	155.0±8.75	53.9±12.56	22.33 ± 4.474	83.7±12.25			
70-74	Male	1799	155.5±8.74	54.5±12.39	22.50 ± 4.640	84.1±12.52			
	Female	1730	155.4±8.55	55.0±12.59	22.70±4.634	84.3±12.32			
75–79	Male	1062	155.1±8.83	53.9±12.83	22.31±4.757	83.5±12.31			
	Female	1061	155.4±8.86	53.9±12.36	22.25 ± 4.458	83.6±12.34			
80-84	Male	618	154.6±8.59	53.9±11.89	22.53±4.529	84.0±11.77			
	Female	605	154.8±8.47	54.5±13.41	22.64 ± 4.840	84.2±13.04			
85–89	Male	276	154.8±7.98	53.6±12.25	22.30±4.439	83.9±12.84			
	Female	299	156.4±9.08	55.8±13.16	22.71±4.499	84.7±13.11			

Data presented as AM±SD. BH: Body height, BW: Body weight, BMI: Body mass index; WC: Waist circumference, AM: Arithmetic mean, SD: Standard deviation

profile of the healthy individuals included in the study with age and sex specifications is shown in Table 1.

The distribution of HGS (kg) in the healthy Indian individuals with age- and sex-wise categorisation is shown in Table 2.

The age- and sex-wise percentile distribution of HGS (kg) in the study population is shown in Table 3.

The age- and sex-wise generated cutoff values of HGS (kg) for the healthy Indian individuals aged 45–89 years are shown in Figure 1.

DISCUSSION

HGS serves as a useful indicator of muscle mass declination and helps in the early detection of sarcopenia.^[2] In the present study, on the basis of the AWGS procedure for determining HGS cutoff values, we propose, a cutoff value of HGS is <15.25 kg for both males and females, for screening, which is an early and relatively easier initial detection of muscle strength deterioration. The value proposed in the present study is similar to the study carried out on Indian older adults by Muhammad *et al.*, where they suggested an HGS cutoff value of 19.5 kg for males and 12.5 kg for females.^[20] The trend continues with female older adults, as another study on Indian adult females reported a value of <16.7 kg; but for males, they reported a relatively higher value of <25.63kg.^[21]

population.						
Age groups (years)	Male	Female				
	HGS (kg)*	HGS (kg)*				
45-49	21.9±7.35	21.9±7.53				
50-54	21.9±7.67	21.7±7.39				
55-59	21.4±7.14	21.5±7.29				
60-64	21.6±7.29	22.0±7.45				
65–69	21.5±7.28	21.5±7.54				
70-74	21.8±7.49	21.9±7.39				
75–79	21.5±7.42	21.8±7.51				
80-84	21.5±6.92	21.8±7.38				
85-89	21.4 ± 7.30	22.3 ± 7.72				

Table 2: The distribution of handgrip strength (kg) in the study

*Presented in AM±SD form. HGS: Handgrip strength, AM: Arithmetic mean, SD: Standard deviation

Looking at the findings from other countries, it is found that the value proposed in the present study has a similarity with a study carried out with older adults of Taiwan; it proposed a value of 19.73 kg.^[22] However, it should also be mentioned that the value proposed in the present study is little less compared to values reported by Chinese (28.5 for males and 18.6 kg for females), Japanese (28 kg for males and 18 kg for females) and Korean (<28.9 kg for males and <16.8 kg

Table 3: Age- and sex-wise percentile distribution of HGS (kg) among the study population.									
Age groups (years)	Sex	Percentiles of HGS (kg)							
		10 th	20 th	25 th	50 th	75 th	90 th	95 th	
45-49	Male	13.1	15.6	16.6	21.1	26.6	32.3	35.5	
	Female	12.7	15.4	16.5	21.1	26.9	32.6	35.4	
50-54	Male	12.5	15.3	16.3	20.9	27.0	32.9	35.7	
	Female	12.6	15.1	16.1	21.0	26.5	32.0	35.5	
55–59	Male	12.8	15.1	16.3	20.6	26.0	31.8	34.5	
	Female	12.8	15.3	16.3	20.9	26.0	32.1	34.9	
60–64	Male	12.9	15.3	16.5	21.0	26.4	31.9	35.2	
	Female	12.9	15.5	16.5	21.3	27.0	32.5	35.4	
65–69	Male	12.8	15.0	16.0	20.6	26.3	31.9	34.8	
	Female	12.4	15.0	16.0	20.5	26.4	32.4	35.4	
70–74	Male	12.5	15.3	16.3	21.3	26.5	32.4	35.8	
	Female	12.8	15.4	16.5	21.0	26.8	32.4	35.4	
75–79	Male	12.8	14.9	16.0	20.5	26.5	32.5	34.8	
	Female	12.9	15.1	16.1	21.0	26.7	32.4	35.6	
80-84	Male	13.0	15.3	16.3	21.0	25.9	31.5	34.3	
	Female	12.9	15.4	16.4	20.8	26.5	32.4	35.5	
85–89	Male	12.8	15.0	15.9	20.1	26.3	32.0	34.9	
	Female	12.8	15.3	16.1	22.0	27.9	32.9	35.1	

HGS: Handgrip strength



Figure 1: Age-specified sex-wise handgrip strength (kg) cutoff values for both Indian (a) males and (b) females (using lowest 20th percentile). HGS: Handgrip strength.

for females) older populations, depicted the prevailing differences in HGS among the Asian populations.^[17,23,24] Our developed cutoff value of HGS of 15.25 kg is quite less than the cutoff provided by AWGS (<28 kg for men and <18 kg for women for Asian population in general) and EWGSOP2 (<27kg for men and <16 kg for women for Europeans).

Limitations

There are several limitations in the study, such as; the data used in the study are cross-sectional in nature, and the study did not consider the nutritional status or physical performance of the individuals for generation of HGS cutoffs; also the determination of handedness of the respondents was not mentioned.

Another important drawback is the study consists of Indians varying with respect to ethnicity and age differences.

However, the study can glimpse the HGS cutoff values for older Indian adults with a large dataset and AWGS procedure, which is worthy of first in its category. The study also reveals the need for gold standard HGS cutoff values with global standardisation for older adults.

CONCLUSION

The study is able to provide a cutoff value of 15.25 kg for older Indians from a large dataset following the AWGS procedure and could be the very first such attempt for older Indians. It implies that individuals with measured HGS values of \leq 15.25 kg run a risk of muscle strength deterioration that may lead to sarcopenia in the long run having serious implications. The outcome of the present work may hence facilitate a screening, that is an early, easy, and cost-effective detection of sarcopenia in terms of HGS values.

Acknowledgements: The authors acknowledge IIPS, Mumbai, India for providing the dataset of LASI, Wave I.

Data availability: The data were freely available in the public domain and can be accessed from IIPS, on subsequent request through https://www.iipsindia.ac.in/content/LASI-data.

Ethical approval: The Institutional Review Board approval is not required.

Declaration of patient consent: Patient's consent was not required as there are no patients in this study.

Financial support and sponsorship: Nil.

Conflicts of interest: There are no conflicts of interest.

Use of artificial intelligence (AI)-assisted technology for manuscript preparation: The authors confirm that there was no use of artificial intelligence (AI)-assisted technology for assisting in the writing or editing of the manuscript and no images were manipulated using AI.

REFERENCES

- 1. Petermann-Rocha F, Chen M, Gray SR, Ho FK, Pell JP, Celis-Morales C. New versus old guidelines for sarcopenia classification: What is the impact on prevalence and health outcomes? Age Ageing 2020;49:300-4.
- 2. Luengpradidgun L, Chamroonkul N, Sripongpun P, Kaewdech A, Tanutit P, Ina N, *et al.* Utility of handgrip strength (HGS) and bioelectrical impedance analysis (BIA) in the diagnosis of sarcopenia in cirrhotic patients. BMC Gastroenterol 2022;22:159.
- 3. Lee SY. Handgrip strength: An irreplaceable indicator of muscle function. Ann Rehabil Med 2021;45:167-9.
- 4. Halaweh H. Correlation between health-related quality of life and hand grip strength among older adults. Exp Aging Res

2020;46:178-91.

- Santra T, Chatterjee S, Chatterjee S, Bardhan S, Banerjee N, Pal B, *et al.* Assessing Lung functions status in male human resources engaged in wood processing works: Using surrogate markers. In: Chakrabarti D, Karmakar S, Salve UR, editors. Ergonomics for design and innovation. Singapore: Springer Nature; 2022. p. 1423-33.
- Mondal S, Bardhan S, Chatterjee S, Chatterjee S, Banerjee N, Mukherjee S. Relationship between body balance and nutritional status in Indian elderly. Indian J Gerontol 2024;38:384-94.
- Neves T, Fett CA, Ferriolli E, Souza MG, Dos Reis Filho AD, Lopes MB, *et al.* Correlation between muscle mass, nutritional status and physical performance of elderly people. Osteoporos Sarcopenia 2018;4:145-9.
- 8. Chatterjee S, Banerjee N, Bhattacharjee S, Santra T, Chatterjee A, Chatterjee S, *et al.* Association of muscular strength and flexibility with select anthropometric markers of obesity: A study on adult Bengalee Kathak danseuses. Int J Curr Adv Res 2017;6:4310-5.
- Yoo JI, Choi H, Ha YC. Mean hand grip strength and cut-off value for Sarcopenia in Korean adults using KNHANES VI. J Korean Med Sci 2017;32:868-72.
- Marques LP, Confortin SC, Ono LM, Barbosa AR, D'Orsi E. Quality of life associated with handgrip strength and sarcopenia: EpiFloripa aging study. Arch Gerontol Geriatr 2019;81:234-9.
- 11. Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyere O, Cederholm T, *et al.* Sarcopenia: Revised European consensus on definition and diagnosis. Age Ageing 2019;48:16-31.
- 12. Chen LK, Woo J, Assantachai P, Auyeung TW, Chou MY, Iijima K, *et al.* Asian working group for sarcopenia: 2019 consensus update on sarcopenia diagnosis and treatment. J Am Med Dir Assoc 2020;21:300-7.e2.
- 13. International Institute for Population Sciences (IIPS), National Programme for Health Care of Elderly (NPHCE), MoHFW, Harvard T. H. Chan School of Public Health (HSPH) and the University of Southern California (USC). Longitudinal ageing study in India (LASI) Wave 1, 2017-18, India report. Mumbai: International Institute for Population Sciences; 2020.
- 14. Dubey AK, Sahoo J, Vairappan B, Parameswaran S, Ps P. Prevalence and determinants of sarcopenia in Indian patients with chronic kidney disease stage 3 and 4. Osteoporos Sarcopenia 2021;7:153-8.
- 15. De A, Kumari S, Kaur A, Singh A, Kalra N, Singh V. Handgrip strength as a screening tool for sarcopenia in males with decompensated cirrhosis. Indian J Gastroenterol 2022;41:284-91.
- Pai M, Muhammad T. Examining factors contributing to the socioeconomic inequalities in handgrip strength among older adults in India: A decomposition analysis. Sci Rep 2024;14:442.
- 17. Kim CR, Jeon YJ, Kim MC, Jeong T, Koo WR. Reference values for hand grip strength in the South Korean population. PLoS One 2018;13:e0195485.
- Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, *et al.* Sarcopenia: European consensus on definition and diagnosis: Report of the European working group on sarcopenia in older people. Age Ageing 2010;39:412-23.

- Chen LK, Liu LK, Woo J, Assantachai P, Auyeung TW, Bahyah KS, et al. Sarcopenia in Asia: Consensus report of the Asian working group for sarcopenia. J Am Med Dir Assoc 2014;15:95-101.
- 20. Muhammad T, Hossain B, Das A, Rashid M. Relationship between handgrip strength and self-reported functional difficulties among older Indian adults: The role of self-rated health. Exp Gerontol 2022;165:111833.
- Singh SA, Madan K, Prabhudesai AA, Agarwal AR, Rastogi R, Bhargava R, *et al.* Normative cutoffs of muscle mass, muscle strength, and frailty for healthy Indian population. Indian J Gastroenterol 2022;41:591-8.
- 22. Lin YH, Chen HC, Hsu NW, Chou P. Using hand grip strength to detect slow walking speed in older adults: The Yilan study.

BMC Geriatr 2021;21:428.

- 23. Ge S, Du Q, Feng X, Liu Y, Wang H, Hai S, *et al.* Optimal cutoffs for the diagnosis of sarcopenia in older Chinese adults. Front Nutr 2022;9:853323.
- Seino S, Kitamura A, Abe T, Taniguchi Y, Murayama H, Amano H, *et al.* Dose-response relationships of sarcopenia parameters with incident disability and mortality in older Japanese adults. J Cachexia Sarcopenia Muscle 2022;13:932-44.

How to cite this article: Mondal S, Banerjee S, Mukherjee S. Handgrip strength cut off value estimation in Indian older adults using LASI-1 dataset. Indian J Physiol Pharmacol. doi: 10.25259/IJPP_585_2024